



REVIEW

Expert Review of the Evidence Base for Arthropod Bite Avoidance

Larry I. Goodyer, PhD,* Ashley M. Croft, MD,[†] Steve P. Frances, PhD,[‡] Nigel Hill, PhD,[§] Sarah J. Moore, PhD,^{§||} Sangoro P. Onyango, BSc,^{§||} and Mustapha Debboun, PhD[¶]

*Leicester School of Pharmacy, De Montfort University, Gateway, Leicester, UK; [†]Surgeon General's Department, London, UK; [‡]Australian Army Malaria Institute, Brisbane, Qld., Australia; [§]London School of Hygiene and Tropical Medicine, London, UK; ^{||}Ifakara Health Institute, Ifakara, Morogoro, Tanzania; [¶]US Army Medical Department Center & School, Center for Health Education & Training, Fort Sam, Houston, TX, USA

DOI: 10.1111/j.1708-8305.2010.00402.x

Bite avoidance measures are commonly recommended to international travelers to help reduce the risk of various arthropod-borne diseases. A key strategy is the use of repellents applied topically to skin or clothing which are considered in the first part of this review. Also advised are a variety of methods that employ the use of insecticides and physical barriers such as mosquito nets or oil preparations applied to the skin.

In the following document, the authors considered some of the most widely used bite avoidance methods and identified the strength and quality of evidence that determined efficacy. The overall purpose of the review is to provide the available evidence, in a graded format, upon which to base recommendations for the selection of appropriate repellents and other methods of bite avoidance in those traveling overseas.

Desired Outcomes and Methods

The authors were asked to consider the effectiveness of the most commonly used active ingredients (AIs) in repellent formulations and methods of bite avoidance. The evidence base considered protection against nuisance biting insects, reduction in the incidence of arthropod-borne diseases, and safety profile. Effectiveness of the repellent related to spectrum of activity against various mosquito species and other arthropods was examined as well as longevity of applied dose. Where possible, efficacy was compared to deet as being the accepted gold standard. All sections employed MEDLINE via PubMed in literature searches augmented by others depending on the subject area investigated. Details of the review process can be found

at www.istm.org; click on "ISTM Committees" and then "Publications."

Part 1: Repellents for Topical Use

Major Findings

N,N-diethyl-3-methylbenzamide (deet), (2-(2-hydroxyethyl)-piperidinecarboxylic acid 1-methyl ester (icaridin), *p*-methane 3, 8-diol (PMD), and ethyl butylacetylaminopropionate (IR3535)-based repellents all provide protection against biting arthropods, but volatile oils and other natural products are less reliable. On the strength of available evidence, the first-line choice for those visiting areas where malaria or other arthropod-borne diseases are endemic remains formulations with higher concentrations (20–50%) of deet. Higher concentration icaridin and PMD preparations are the most useful alternatives to deet where they are available. See Table 1 for a summary of the findings.

Diethyltoluamide (Other Name Deet; Chemical Name: *N,N*-Diethyl-3-Methylbenzamide, Former Nomenclature *N,N*-Diethyl-*m*-Toluamide)

Deet has been widely used in insect repellent products for use on human skin to protect against biting arthropods.^{1–5} The majority of laboratory and field tests conducted to compare the efficacy of repellents use deet as the primary formulation or as a comparison. Deet is considered the most effective broad spectrum repellent AI against biting arthropods.⁶

The first laboratory tests against mosquitoes were reported by Gilbert and colleagues⁷ who showed deet and dimethylphthalate were equally effective against *Anopheles quadrimaculatus*. Altman⁸ reported field studies in Panama against *Anopheles albimanus* and showed 75% deet provided protection for at least 3 hours. Field studies undertaken in the last 20 years in Africa,^{9,10} Australia,^{11,12} Papua New Guinea,^{13,14} Malaysia,¹⁵ and Thailand¹⁶ have shown that protection

Corresponding Author: Larry I. Goodyer, PhD, Leicester School of Pharmacy, DeMontfort University, The Hawthorn Building, Gateway, Leicester LE19BH, UK. E-mail: LGoodyer@dmu.ac.uk

Table 1 Summary of evidence base for topically applied repellents

Aspect of bite avoidance	Strength of evidence*	Quality of evidence†	Comments
Deet			
Dermal application to avoid mosquitoes	A	I	Deet provides shorter protection against <i>Anopheles</i> sp. mosquitoes than Culicine mosquitoes. Reapplication times will vary to maintain optimal effectiveness.
Dermal application to avoid ticks	B	I	Laboratory and field tests showed deet provides minimal protection, recent test showed adequate protection.
Application of deet to wristbands	E	I	Wristbands provide no protection for uncovered and untreated human skin.
Icaridin			
Dermal application to avoid mosquitoes	A	I	Icaridin provided good protection against <i>Anopheles</i> sp. mosquitoes in Africa and Asia, also effective against Culicine mosquitoes.
Dermal application of icaridin to humans to avoid ticks	B	I	Laboratory test showed icaridin provides protection for 1 h. Recent test showed adequate protection.
IR3535			
Dermal application to avoid mosquitoes	B	II	Based on limited field studies for mosquitoes. Not recommended in malaria endemic areas.
Dermal application to avoid sand flies	B	II	Based on one laboratory study.
PMD			
Dermal application to avoid mosquitoes	A	I	Can be highly recommended as an alternative repellent to deet at concentrations of >20% as a repellent for use in disease endemic areas. (Some evidence of efficacy against ticks.)
Citronella	C	III	Not recommended for use when engaging in vigorous activities, in disease endemic areas or areas with high densities of mosquitoes.
Neem	D	III	More studies should be conducted before it is recommended as a repellent.
Essential oils	D	III	Essential oils require careful formulation to be effective, and safety data suggest skin irritation is a factor.

*A = good evidence to support use; B = moderate evidence to support use; C = poor evidence to support use; D = moderate evidence to support a recommendation against use; E = good evidence to support a recommendation against use.

†I = evidence from one or more properly randomized, controlled trial; II = evidence from one or more well-designed clinical trial without randomization, from case-controlled analysis of cohort study; III = consensus evidence, evidence from one authority or reports of expert committees, single case studies.

against *Anopheles* spp. is less than that provided against Culicine mosquitoes.

The response of different mosquito species to deet is variable.¹⁷ Field tests of repellent formulations containing deet against biting *Culex* spp., *Aedes* spp., *Mansonia* spp., and *Verrallina* spp. have been reported.⁵ The protection provided by deet was longer against these genera than provided against *Anopheles* spp.¹²

Studies have shown that deet provides only minimal or poor protection against ticks.^{18–21} However, recently Carroll and colleagues²² showed that a 33% deet, Extended Duration formulation provided high levels of protection for 12 hours.

Deet is recommended to be applied to the exposed skin of humans. However, alternative methods of using deet have been proposed and investigated. The application of deet to wide mesh cotton/nylon jackets provided good protection against mosquitoes and biting flies.²³ Deet-treated netting used as groundsheets were shown to provide significant protection against ticks.²⁴ Although application of deet to nylon/cotton fabrics has been shown to enhance protection against bites, the application of deet to some synthetic fibers and plastics

may cause damage, and thus the use of deet applied to clothing is not widely accepted. The use of wristbands treated with deet and other AIs offered no protection against mosquitoes.⁴

There have been a number of reviews concerning the safety of deet,^{25,26} and they have attested to its generally acceptable safety profile. There are few reports of systemic toxicity in adults following dermal application. The safety profile in the second and third trimester of pregnancy has been established through observation of very low placental cord concentrations after maternal application of deet,²⁷ and animal models do not indicate any teratogenic effects.²⁸ Recommendations for use in young children do vary between countries, with some recommending lower concentrations²⁹ and others suggesting that higher strengths can be used.³⁰ However, the causation between the few reported cases of encephalopathy in children and the topical use of deet cannot be supported by a good evidence base.³¹

The scientific evidence and continued use of deet for >50 years has shown this AI is the best broad spectrum repellent available for minimizing bites of mosquitoes, ticks, and other biting arthropods.

Icaridin (Formerly Picaridin; (2-(2-Hydroxyethyl)-Piperidinecarboxylic Acid 1-Methyl Ester; WHO Designation: Icaridin; Trade Name: Bayrepel; Development Reference Code: KBR 3023)

Independent field studies demonstrating the effectiveness of repellents containing icaridin against mosquitoes have been conducted in Malaysia^{32,33} and Florida.³⁴ In Australia, a formulation containing 19.2% icaridin provided similar protection as 20% deet against *Verrellina lineata*.³⁵ In another study in Australia, the same formulation provided >95% protection against *Culex annulirostris* for 5 hours, but only 1 hour protection against *Anopheles* spp.¹² KBR 3023 at concentrations of 2% to 13% v/v in 90% ethanol provided better protection against Anophelines in Africa than comparable formulations containing deet.¹⁰

Field studies against mosquitoes in two locations in Australia showed that a 9.3% formulation only provided 2-hour protection against *V. lineata*³⁵ and 5-hour protection against *C. annulirostris*,³⁶ while 7% icaridin provided 5.7 hours of protection against *Aedes albopictus* in laboratory tests.³⁷ The use of lower concentrations of icaridin in commercial formulations may require the user to reapply repellent more often to maintain effectiveness than with the higher concentrations (>20%) of icaridin used in the field.

Protection from biting by ticks provided by 20% lotions of KBR 3023 was reported to be short.³⁸ Carroll and colleagues²² showed that Bayrepel (10 and 20% icaridin) repellent provided high levels of protection for 12 hours when applied to human volunteers against *Amblyomma americanum* under simulated field-contact conditions.

IR3535 [Also known as 3535 or EBAAP (Ethyl Butylacetylaminopropionate)]

Five field studies were identified, all testing IR3535 against mosquitoes.^{10,34,39–41} These indicated that IR3535 is as effective as deet in repelling mosquitoes of the *Aedes* and *Culex* genera but may be less effective than deet in repelling anopheline mosquitoes. A number of laboratory studies were also identified, testing IR3535 against a variety of other arthropods, including blackflies and ticks.⁴² An uncontrolled field study of a new, controlled-release formulation of IR3535 reported that these formulations may provide complete protection against mosquito biting for 7.1 to 10.3 hours.⁴¹ IR3535 may be more effective than deet in protecting against phlebotomine sandfly biting (10.4 h mean protection vs 8.8 h, respectively).⁴²

PMD: Lemon Eucalyptus (Corymbia citriodora) Extract

The principal repellent component of lemon eucalyptus extract is PMD, which is the main by-product of lemon eucalyptus hydrodistillation.⁴³ The active component is prepared through acid modified extraction of leaves or a synthetic version of PMD is used in the majority of

commercially available preparations. Importantly, PMD has been proven to prevent malaria in a clinical trial in the Bolivian Amazon.⁴⁴

Studies carried out both in the laboratory and the field using rigorous methodology have shown PMD to be a repellent of equal efficacy and longevity as deet.⁴⁵ At 30% AI, PMD provided almost complete protection for 4 hours in South America⁴⁶ and complete protection for 6 hours at 50% AI in Sub-Saharan Africa against malaria vectors.⁴⁷ In both of these studies, the protection time was equivalent to the deet controls. A well-designed laboratory trial of PMD against a further African malaria vector showed complete protection for 4 to 5 hours using PMD impregnated towlettes,⁴⁸ again comparable with deet. Laboratory trials using the main vectors of dengue fever have shown good protection, which is important for travelers as the vector bites in the day-time.^{45,49} Against the tick vectors of Lyme disease and Rocky Mountain spotted fever, PMD reduces attachment and feeding success by around 77%, and PMD is highly effective against the Highland Midge.⁵⁰ PMD has not been tested against the vectors of leishmaniasis in vivo, although in vitro results suggest that it may be effective.⁵¹

Citronella

Citronella is one of the essential oils obtained from the leaves and stems of different species of *Cymbopogon* grasses. From the available literature and information, we can conclude that the complete protection time for citronella-based repellents is <2 hours^{4,49,52} because the repellent is highly volatile, but this can be prolonged by careful formulation and the addition of fixatives like vanillin.⁵³

Neem

Neem is a vegetable oil pressed from the fruits and seeds of neem (*Azadirachta indica*). Several field studies from India have shown very high efficacy of neem-based preparations.^{54–56} However, these studies have used questionable methodologies and their results contrast strongly with several others that have shown medium-range protection from neem products being inferior to deet.^{46,49,57} Neem has a low dermal toxicity but can cause skin irritation such as dermatitis.⁵⁸ However, caution should be taken as neem is a proven reproductive toxicant and long-term subchronic exposure could impair fertility.⁵⁹

Essential Oils

Many commercial repellents contain a number of plant essential oils either for fragrance or as repellents. The most effective of these include thyme oil, geraniol, peppermint oil, cedar oil, patchouli, and clove.^{52,60,61} Most of these essential oils are highly volatile and this contributes to their poor longevity as mosquito repellents. They can be irritating to the skin^{49,62} and their repellent effect is variable, dependent on formulation and concentration.

Conclusions

The largest body of evidence for effectiveness in terms of spectrum of activity and longevity relates to deet that remains as a gold standard to which newer repellents are compared in reducing nuisance bites from arthropods. Icaridin and PMD are reasonable alternatives to deet for those visiting areas where arthropod-borne diseases are endemic, whereas IR3535 has shown reduced efficacy against *Anopheles* mosquitoes and should not be advised for malaria endemic areas. When advising a formulation, the concentration of AI and the expected application rate of AI should always be considered because these will greatly influence longevity of the applied dose. There are, for instance, some icaridin formulations containing suboptimal concentrations. Apart from the repellent choice, the following factors will determine the duration of any repellent product.

1. Product concentration: In general, the higher percentage of AI, the greater the protection time will be, although this tends to plateau at 50% w/v in the case of deet.⁶³
2. The rate of application that is applied to the exposed human skin is also variable.⁶⁴ And travelers have been shown to apply relatively low doses of AI on treated skin.⁶⁵
3. Activity level of the user: The effectiveness of a repellent is reduced by evaporation and absorption from the skin surface and wash-off by sweat.
4. Environmental conditions: Rain or water, washing the repellent off the body, wind, and high ambient temperatures will reduce a product's effectiveness.
5. User's overall attractiveness to blood-feeding arthropods, such as mosquitoes.
6. Number and species of blood-feeding arthropods (biting density).

Part 2: Area and Barrier Methods

Major Findings

The strongest level of evidence exists for the use of insecticide-treated mosquito nets, and these are to be advised for all travelers visiting disease endemic areas at risk from biting arthropods on retiring. Insecticide-treated clothing and other fabrics would also be a useful adjunct to dermal applied repellents. Electric insecticide vaporizers, essential oil candle, and coils to burn do reduce bites from arthropods, but there is little evidence on the efficacy of knockdown insecticide sprays. There is some concern regarding the potential adverse effects of burning coils. There is less evidence that these technologies reduce the incidence of malaria. There is only weak evidence regarding the efficacy of oils used on the skin. See Table 2 for a summary of the findings.

Fabric Impregnation With Insecticides

The use of fabric impregnated with insecticides, particularly insecticide-treated bed nets, has become an

important tool or method of personal protection against arthropod bites and disease-transmitting vectors. Some of the insecticides that are recommended and used for treatment of fabrics are permethrin, deltamethrin, lambda-cyhalothrin, alpha-cypermethrin, cyfluthrin, and etofenprox.⁶⁶ However, the insecticide most commonly used for fabric impregnation is permethrin [3-(phenoxyphenyl) methyl (±)-*cis*, *trans*-3-(2,2-dichloroethenyl)-2,2-dimethyl-cyclopropanecarboxylate].

Permethrin is a synthetic pyrethroid insecticide derived from crushed dried flowers of the plant *Chrysanthemum cinerarifolium*. Although permethrin's primary mode of action is contact toxicity against a wide variety of biting arthropods, it is also unique in that it serves both as a contact insecticide and as an insect repellent. Permethrin-impregnated clothing provides good protection against mosquitoes,^{67–77} ticks,^{78–84} chigger mites,^{85,86} fleas,⁸⁷ lice,^{88,89} sand flies,^{90,91} kissing bugs,^{92,93} and tsetse flies.⁹⁴ Thus, the use of permethrin-treated clothing will decrease the biting frequency and transmission of arthropod-borne diseases among civilian travelers and deployed military personnel. Today, military personnel from many countries use permethrin to repel and kill arthropods that land on many kinds of treated surfaces, including field uniforms, tents, bed nets, and helmet covers.⁹⁵

Impregnated-treated fabrics such as bed nets, curtains, chaddars (veils or wraps worn by Muslim women), top sheets, and blankets have also been found to be effective in reducing the burden of malaria and other vector-borne diseases^{96–100} and have been used in the Roll Back Malaria Program by the World Health Organization for tropical countries. However, due to the development of pyrethroid resistance in mosquito vectors, the use of impregnating fabrics with insecticide mixtures, a combination of a repellent and insecticide-treated bed nets or mixtures of repellents and non-pyrethroid-treated fabrics have become new promising tools for disease vector control.^{101,102}

The development and use of long-lasting insecticide nets such as Olyset Net and Perma Net has also been proven to be effective in reducing or eliminating the need for insecticide retreatment of insecticide-treated nets.^{103–107} In addition, recently, the use of tent barrier treatments with bifenthrin and permethrin is another effective method of personal protection against biting arthropods.^{108–111}

Electric Insecticide Vaporizers

An insecticide vaporizer, which are mains- or battery-operated electrical devices releasing a pyrethroid insecticide, will clear a room of insects in around 30 minutes and will remain effective for over 6 hours.¹¹²

Ten field and laboratory studies were identified, testing a variety of insecticides and devices in different settings, against a range of flying insects, including various mosquito species.^{113–122}

Table 2 Summary of evidence base for area and barrier methods

Effect	Strength of evidence*	Quality of evidence†	Comments
Insecticides			
Effectiveness of insecticides on clothing	A	I	Strong evidence from numerous authorities or reports of expert committees supporting the use of impregnated insecticides (permethrin) on clothing as an important method of personal protection against arthropod bites.
Effectiveness of net impregnation	A	I	Numerous studies have shown good personal protection against arthropod bites with the use of nets impregnated with insecticides.
Effectiveness of use on tents	B	II	Consistent evidence from studies showing good personal protection against arthropod bites with the use of impregnated tents with insecticides.
Vaporizing insecticide mats			
Inhibition of nuisance biting by mosquitoes	B	II	Consistent evidence that insecticide vaporizers cause mosquito bite inhibition, mosquito repellence, mosquito knockdown, and mosquito death.
Malaria prevention	D	II	No evidence from three studies
Mosquito coils			
Inhibition of nuisance biting by mosquitoes	B	II	Good evidence of protection against mosquito bites
Malaria prevention	C	III	Consensus that mosquito coils could be a useful measure to prevent malaria
Lung cancer	D	II	Prior exposure to mosquito coil smoke was more common in lung cancer patients than in controls. Lung cancer risk was higher in frequent burners of mosquito coils than in non-burners. ¹²⁵
Other adverse effects	D	III	Breathlessness, ¹³² chest irritation, ¹¹⁴ eye and nose irritation. ^{120,132,147}
Essential oil candles			
Inhibition of nuisance biting by mosquitoes	C	II	Two studies demonstrating significant repellency compared to control. ^{73,134}
Inhibition of nuisance biting by other insect species	C	II	Essential oil, linalool, and geraniol but not citronella candles repelled significantly more sandflies than did the control candle. ¹³⁵
Mosquitoes knockdown sprays			
Inhibition of nuisance biting by mosquitoes and other insect species	C	III	Anecdotal evidence only.
Malaria prevention	D	II	Two studies showed no significant reduction in malaria incidence. ^{119,137}
Prevention of other insect-transmitted infections	D	III	No studies.
Oil-based products			
Inhibition of nuisance biting by midges	C	I	Evidence for bath oils only.
Inhibition of nuisance biting by mosquitoes	C	II	Evidence for chemical base oils only. Inhibition was observed mostly in culicine mosquitoes. ¹⁴⁸
Soybean oil inhibition of nuisance biting by mosquitoes and black fly	D	III	More studies (field trials) needed to establish the efficacy of soybean oil as a repellent, although initial studies are very promising. Toxicity very low.
Garlic and vitamin B	E	II	Do not confer protection against haematophagous arthropods and this notion should be discouraged.

*A = good evidence to support use; B = moderate evidence to support use; C = poor evidence to support use; D = moderate evidence to support a recommendation against use; E = good evidence to support a recommendation against use.

†I = evidence from one or more properly randomized, controlled trial; II = evidence from one or more well-designed clinical trial without randomization, from case-controlled analysis of cohort study; III = consensus evidence, evidence from one authority or reports of expert committees, single case studies.

Electric insecticide vaporizers using pyrethrins or pyrethroids inhibit nuisance biting by mosquitoes, and vaporized pyrethrins kill house flies. There is no direct evidence that the technology does prevent malaria or any other insect-transmitted infection.^{119,120,122} Exposure to pyrethrins and/or pyrethroids may present a low-level health hazard to humans,¹²³ and so

vaporizers containing these classes of insecticide should be used by travelers with caution. Vaporized neem oil is an effective inhibitor of mosquito landing and appears safe to humans.¹²¹ It should be investigated further for its potential usefulness in preventing malaria and other mosquito-transmitted infections.

Mosquito Coils

Mosquito coils are made from a paste of powdered insecticide, usually a pyrethrin or pyrethroid which when lit will smolder at a steady rate. Burning one mosquito coil releases the same amount of particulate matter as does burning 75–137 cigarettes¹²⁴ and emits formaldehyde equivalent to 51 cigarettes.¹²⁵

Historically, some authorities have advised international travelers to burn mosquito coils in their room at night, in order to discourage mosquito biting and so help prevent mosquito-transmitted infection, notably malaria. These authorities include some national expert bodies.^{126–130}

Consistent evidence from a systematic review indicated that coils cause mosquito bite inhibition, mosquito repellence, mosquito deterrence, mosquito knockdown, and mosquito death. Inhibition of nuisance biting is seen in all mosquito species: *Aedes*, *Anopheles*, *Culex*, and *Mansonia*.¹³¹ There has been little evidence that this technology prevents malaria transmission^{120,131,132} or other mosquito-transmitted infection. However, the expert group considers that they would be effective in preventing malaria in the light of as yet unpublished data.¹³³ Exposure to mosquito coil smoke could be hazardous to humans due to its potential association with lung cancer, but this should be taken in context of the amount likely to be inhaled over an evening and the very occasional use by travelers. In many countries, such coils are licensed for outdoor use only due to these concerns.

Essential Oil Candles

Three field studies were identified, demonstrating the effectiveness of essential oil candles in repelling mosquitoes and sand flies.^{134–136}

Burning essential oil candles is likely to prevent biting by both mosquitoes and by sandflies. They may also prevent biting by other insect species. While there is no evidence that this technology prevents malaria, leishmaniasis, or any other insect-transmitted disease, this is an aspect which should be investigated. Candles containing 5% essential oil of geraniol appear to hold the most promise.

Knockdown Insecticide Sprays

Knockdown insecticides are aerosol sprays which are designed to be sprayed indoors and into the air, to eliminate flying insects by killing them as they fly through the room.¹²⁸

Two individual studies were identified which failed to demonstrate that knockdown insecticide sprays prevented malaria in travelers to Africa.^{119,132} Only anecdotal evidence supports the assumption that knockdown sprays inhibit nuisance biting by flying insects. There is an obvious, but mostly unquantified health risk to humans, from inhaling any insecticide vapor.¹³⁷ In the absence of persuasive evidence on the benefits of this technology, the use of knockdown

insecticide sprays should be discouraged, in favor of vector avoidance strategies of proven effectiveness.¹³⁸

Bath Oils and Chemical Base Oils

Bath oils, and chemical base oils also, seem to protect against insect biting not by a repellent action but by forming a physical barrier between the human target and the insect.¹³⁹ They are reported to be especially effective against small flying insects, creating an oily layer which traps these insects on the sticky surface of the skin.¹⁴⁰ Some studies have suggested that small flying insects, such as biting midges and sandflies, are not efficiently repelled by conventional repellants (deet and pyrethroid insecticides).^{141,142}

One small randomized controlled trial (nine adult volunteers) tested a commercial bath oil preparation (Avon Skin-so-Soft, SSS)¹⁴⁰ and found that deet formulations were significantly more effective in preventing midge biting than was SSS.

Soybean Oil

Two well-designed laboratory evaluations of Bite Blocker, a commercial preparation containing 2% soybean oil in addition to other oils and emulsifiers, have shown that it is competitive with deet, against a dengue vector and nuisance biting mosquitoes in one study⁴⁹ and equivalent to that of low-concentration deet in a second study.⁴ A field trial showed 3.5-hour protection under intensive biting pressure of nuisance mosquitoes, but this was not conducted by independent researchers.¹⁴³ In a similar study against black flies, soybean oil provided complete protection from black fly bites of 9.7 hours as compared to 6.6-hour protection provided by deet.¹⁴⁴

Garlic and Vitamin B

It is still commonly believed and reported in magazines that ingestion of garlic or B vitamins makes the human skin unpalatable to blood feeding and biting arthropods which have been refuted.^{145,146} Garlic and B vitamins must never be suggested as a natural method of bite prevention.

Conclusions

The use of insecticide-treated mosquito nets and clothing is well supported by the data and is to be recommended to travelers visiting malaria endemic areas. Electric insecticide vaporizers and essential oil candles inhibit nuisance biting, but there is little evidence that they help prevent malaria. Mosquito coils are effective and may help to reduce the risk of malaria, although safety concerns have been raised. The use of bath oils and other oils should be discouraged in travelers until further effective personal protection evidence is available.¹²⁷

Acknowledgment

The authors dedicate this review to the memory of Dr Nigel Hill who died suddenly in January 2010.

Declaration of Interests

L. I. G. is director of Nomad Medical that produces deet and permethrin based products. A. M. C., N. H., S. M., and P. S. state that they have no conflict of interest. The opinions expressed herein are those of the authors and do not necessarily reflect those of the UK Ministry of Defence, the United States Department of Defense, and the Joint Health Command of the Australian Defence Force or any current defense policy.

References

- Curtis CF, Lines JD, Baolin L, Renz A. Natural and synthetic repellents. In: Curtis CF, ed. *Appropriate technology in vector control*. Boca Raton, FL: CRC Press, 1990:75–92.
- Brown H, Hebert AA. Insect repellents: an overview. *J Am Acad Dermatol* 1997; 36:243–249.
- Fradin MS. Mosquitoes and mosquito repellents: a clinician's guide. *Ann Intern Med* 1998; 128:931–940.
- Fradin MS, Day JF. Comparative efficacy of insect repellents against mosquito bites. *N Engl J Med* 2002; 347:13–18.
- Frances SP. Efficacy and safety of repellents containing deet. In: Debboun M, Frances SP, Strickman D, eds. *Insect repellents: principles, methods & uses*. 1st Ed. Boca Raton, FL: CRC Press, 2007:311–325.
- Gupta RK, Rutledge LC. Role of repellents in vector control and disease prevention. *Am J Trop Med Hyg* 1994; 50:82–86.
- Gilbert IH, Gouck HK, Smith CN. New mosquito repellents. *J Econ Entomol* 1955; 48:741–743.
- Altman RM. Repellent tests against *Anopheles albimanus* Wiedemann in the Panama Canal Zone. *Mosq News* 1969; 29:110–112.
- Walker T, Robert LL, Copeland A, et al. Field evaluation of arthropod repellents, deet and a piperidine compound, AI3-37220, against *Anopheles funestus* and *Anopheles arabiensis* in Western Kenya. *J Am Mosq Control Assoc* 1996; 12:172–176.
- Constantini C, Badolo A, Ilboudo-Sanogo E. Field evaluation of the efficacy and persistence of insect repellents deet, IR3535 and KBR 3023 against *Anopheles gambiae* complex and other Afrotropical vector mosquitoes. *Trans R Soc Trop Med Hyg* 2004; 98:644–652.
- Frances SP, Cooper RD, Sweeney AW. Laboratory and field evaluation of the repellents deet, CIC-4, and AI3-37220 against *Anopheles farauti* (Diptera: Culicidae) in Australia. *J Med Entomol* 1998; 35:690–693.
- Frances SP, Waterson DGE, Beebe NW, Cooper RD. Field evaluation of repellent formulations containing deet and picaridin against mosquitoes in Northern Territory, Australia. *J Med Entomol* 2004; 41:414–417.
- Frances SP, Cooper RD, Popat S, Sweeney AW. Field evaluation of the repellents, deet, CIC-4 and AI3-37220, against *Anopheles* (Diptera: Culicidae) in Lae, Papua New Guinea. *J Am Mosq Control Assoc* 1999; 14:339–341.
- Frances SP, Cooper RD, Popat S, Beebe NW. Field evaluation of the repellents containing deet and AI3-37220, against *Anopheles koliensis* in Papua New Guinea. *J Am Mosq Control Assoc* 2001; 17:42–44.
- Yap HH, Jahinger K, Zairi J. Field efficacy of four insect repellent products against vector mosquitoes in a tropical environment. *J Am Mosq Control Assoc* 2000; 16:241–244.
- Frances SP, Klein TA, Hildebrandt DW, et al. Laboratory and field evaluation of the repellents, deet, CIC-4 and AI3-37220, against *Anopheles dirus* (Diptera: Culicidae) in Thailand. *J Med Entomol* 1996; 33:511–515.
- Rutledge LC, Moussa MA, Lowe CA, Sofield RK. Comparative sensitivity of mosquito species and strains to the repellent diethyl toluamide. *J Med Entomol* 1978; 14:536–541.
- Schreck CE, Fish D, McGovern TP. Activity of repellents applied to skin for protection against *Amblyomma americanum* and *Ixodes scapularis* ticks (Acari: Ixodidae). *J Am Mosq Control Assoc* 1995; 11:136–140.
- Salafsky B, He Y-X, Li J, et al. Short report: study on the efficacy of a new long-acting formulation of *N,N*-diethyl-*m*-toluamide (deet) for the prevention of tick attachment. *Am J Trop Med Hyg* 2000; 62:169–172.
- Jensenius M, Pretorius A-M, Clarke F, Myrvang B. Repellent efficacy of four commercial deet lotions against *Amblyomma herbraeum* (Acari: Ixodidae), the principal vector of *Rickettsia africae* in southern Africa. *Trans R Soc Trop Med Hyg* 2005; 99:708–711.
- Solberg VB, Klein TA, McPherson KR, et al. Field evaluation of deet and a piperidine repellent (AI3-37220) against *Amblyomma americanum* (Acari: Ixodidae). *J Med Entomol* 1995; 32:870–875.
- Carroll JF, Benante JP, Klun JA, et al. Twelve-hour duration testing of cream formulations of three repellents against *Amblyomma americanum*. *Med Vet Entomol* 2008; 22:144–151.
- Grothaus RH, Haskins JR, Schreck CE, Gouck HK. Wide mesh netting, and improved method of protection against blood-feeding Diptera. *Mosq News* 1976; 36:11–18.
- Grothaus RH, Reed JT, Passingham LH. Field evaluation of arthropod repellent netting as a ground cloth against ticks. *Am J Trop Med Hyg* 1976; 25:747–750.
- Goodyer L, Behrens RH. Short report: the safety and toxicity of insect repellents. *Am J Trop Med Hyg* 1998; 59:323–324.
- Sudakin DL, Wade R, Trevathan BS. Deet: a review and update of safety and risk in the general population. *J Toxicol Clin Toxicol* 2003; 41:831–839.
- McGready R, Hamilton KA, Simpson JA, et al. Safety of the insect repellent *N,N*-Diethyl-*M*-Toluamide (DEET) in pregnancy. *Am J Trop Med* 2001; 65: 285–289.
- Schoenig GP, Neeper-Bradley TL, Fisher LC, Hartnagel RE Jr. Teratological evaluations of DEET in rats and rabbits. *Fundam Appl Toxicol* 1994; 23:63–9.
- Pest Management Regulatory Agency, Health Canada. Re-evaluation Decision Document RRD2002-01. Available at: <http://www.hc-sc.gc.ca/pmr-arla/english/pdf/rrd/rrd2002-01-e.pdf>. (Accessed 2002 Apr 26).

30. Chiodini P, Hill D, Lalloo D, et al. Guidelines for malaria prevention in travellers from the United Kingdom. London: Health Protection Agency, 2007.
31. Koren G, Matsui D, Bailey B. DEET-based insect repellents: safety implications for children and pregnant and lactating women. *CMAJ* 2003; 169:209–212.
32. Yap HH, Jahangir K, Chong SC, et al. Field efficacy of a new repellent, KBR 3023, against *Aedes albopictus* (Skuse) and *Culex quinquefasciatus* (Say) in a tropical environment. *J Vect Ecol* 1998; 23:62–68.
33. Yap HH, Jahangir K, Zairi J. Field efficacy of four insect repellent products against vector mosquitoes in a tropical environment. *J Am Mosq Control Assoc* 2000; 16:241–244.
34. Barnard DR, Bernier UR, Posey KH, Xue R-D. Repellency of IR3535, KBR3023, *para*-menthane-3,8-diol, and deet to black salt marsh mosquitoes (Diptera: Culicidae) in the Everglades National Park. *J Med Entomol* 2002; 39:895–899.
35. Frances SP, Dung NV, Beebe NW, Debboun M. Field evaluation of repellent formulations against daytime and nighttime biting mosquitoes in a tropical rainforest in northern Australia. *J Med Entomol* 2002; 39:541–544.
36. Frances SP, Waterson DGE, Beebe NW, Cooper RD. Field evaluation of commercial repellent formulations against mosquitoes (Diptera: Culicidae) in Northern Territory, Australia. *J Am Mosq Control Assoc* 2005; 21:480–482.
37. Xue R-D, Ali A, Day JF. Commercially available insect repellents and criteria for their use. In: Debboun M, Frances SP, Strickman D, eds. *Insect repellents: principles, methods & uses*. 1st Ed. Boca Raton, FL: CRC Press, 2007:405–415.
38. Pretorius A-M, Jensenius M, Clarke F, Ringertz SH. Repellent efficacy of deet and KBR 3023 against *Amblyomma hebraeum* (Acari: Ixodidae). *J Med Entomol* 2003; 40:245–248.
39. Thavara U, Tawatsin A, Chompoosri J, et al. Laboratory and field evaluations of the insect repellent 3535 (ethyl butylacetylaminopropionate) and deet against mosquito vectors in Thailand. *J Am Mosq Control Assoc* 2001; 17:190–195.
40. Naucke TJ, Kröpke R, Benner G, et al. Field evaluation of the efficacy of proprietary repellent formulations with IR3535 and picaridin against *Aedes aegypti*. *Parasitol Res* 2007; 101:169–177.
41. Carroll SP. Prolonged efficacy of IR3535 repellents against mosquitoes and blacklegged ticks in North America. *J Med Entomol* 2008; 45:706–714.
42. Naucke TJ, Lorentz S, Grünwald HW. Laboratory testing of the insect repellents IR3535 and deet against *Phlebotomus mascittii* and *P. duboscqi* (Diptera: Psychodidae). *Int J Med Microbiol* 2006; 296(Suppl 40):230–232.
43. Schreck CE, Leonhardt BA. Efficacy assessment of Quwenling, a mosquito repellent from China. *J Am Mosq Control Assoc* 1991; 7:433–436.
44. Hill N, Lenglet A, Arnez AM, Cainero I. Randomised, double-blind control trial of p-menthane diol repellent against malaria in Bolivia. *BMJ* 2007; 335:1023.
45. Carroll SP, Loye J. PMD, a registered botanical mosquito repellent with deet-like efficacy. *J Am Mosq Control Assoc* 2006; 22:507–514.
46. Moore SJ, Lenglet A, Hill N. Field evaluation of three plant-based insect repellents against malaria vectors in Vaca Diez Province, the Bolivian Amazon. *J Am Mosq Control Assoc* 2002; 18:107–110.
47. Trigg JK. Evaluation of a eucalyptus-based repellent against *Anopheles* spp. in Tanzania. *J Am Mosq Control Assoc* 1996; 12:243–246.
48. Govere J, Durrheim DN, Baker L, et al. Efficacy of three insect repellents against the malaria vector *Anopheles arabiensis*. *Med Vet Entomol* 2009; 14:441–444.
49. Barnard DR, Xue RD. Laboratory evaluation of mosquito repellents against *Aedes albopictus*, *Culex nigripalpus*, and *Ochlerotatus triseriatus* (Diptera: Culicidae). *J Med Entomol* 2004; 41:726–730.
50. Trigg JK, Hill N. Laboratory evaluation of a eucalyptus-based repellent against four biting arthropods. *Phytother Res* 1996; 10:313–316.
51. Rowton ED, Grieco JP, Coleman RE. High throughput testing for toxic and behaviour modifying chemicals against Phlebotomine sand flies, DoD Tri-Service Pest Management Workshop; Jacksonville, FL. 2007.
52. Trongtokit Y, Rongsriyam Y, Komalamisra N, Apiwatanasorn C. Comparative repellency of 38 essential oils against mosquito bites. *Phytother Res* 2005; 19:303–309.
53. Tawatsin A, Wratten SD, Scott RR, et al. Repellency of volatile oils from plants against three mosquito vectors. *J Vect Ecol* 2001; 26:76–82.
54. Singh N, Mishra AK, Saxena A. Use of Neem cream as a mosquito repellent in tribal areas of central India. *Indian J Malariol* 1996; 33:99–102.
55. Sharma SK, Dua VK, Sharma VP. Field studies on the mosquito repellent action of Neem oil. *Southeast Asian J Trop Med Public Health* 1995; 26:180–182.
56. Caraballo AJ. Scientific note on the mosquito repellent action of Neemos®. *J Am Mosq Control Assoc* 2000; 16:45–46.
57. Hadis M, Lulu M, Mekonnen Y, Asfaw T. Field trials on the repellent activity of four plant products against mainly *Mansonia* population in western Ethiopia. *Phytother Res* 2003; 17:202–205.
58. Reutemann P, Ehrlich A. Neem oil: an herbal therapy for alopecia causes dermatitis. *Dermatitis* 2008; 19:E12–E15.
59. Boeke SJ, Boersma MG, Alink GM, et al. Safety evaluation of Neem (*Azadirachta indica*) derived pesticides. *J Ethnopharmacol* 2004; 94:25–41.
60. Barnard DR. Repellency of essential oils to mosquitoes (Diptera: Culicidae). *J Med Entomol* 1999; 36:625–629.
61. Rutledge LC, Gupta RK. Reanalysis of the C G Macnay mosquito repellent data. *J Vect Ecol* 1995; 21:132–135.
62. Kiken DA, Cohen DE. Contact dermatitis to botanical extracts. *Am J Contact Dermat* 2002; 13:148–152.
63. Buescher MD, Rutledge LC, Wirtz RA, Nelson JH. The dose-response of deet against *Aedes aegypti*. *Mosq News* 1983; 42:364–366.
64. Gupta RK, Sweeney AW, Rutledge LC, et al. Effectiveness of controlled-release personal-use arthropod repellents and permethrin-impregnated clothing in the field. *J Am Mosq Control Assoc* 1987; 3:556–560.
65. Thrower Y, Goodyer LI. Application of insect repellents by travellers to malaria endemic areas. *J Travel Med* 2006; 13:198–202.
66. WHO. Instructions for treatment and use of insecticide-treated mosquito nets. Geneva: World Health Organization 2000. (WHO/CDS/RBM/2002.41). Available at <http://whqlibdoc.who.int/hq/2002/WHO.CDS.RBM.2002.41.pdf>. (Accessed 2010 Mar).

67. Rozendaal JA. Impregnated mosquito nets and curtains for self-protection and malaria control. *Trop Dis Bull* 1998;86:R1–R41.
68. Schreck CE, Posey K, Smith D. Durability of permethrin as a potential clothing treatment to protect against blood-feeding arthropods. *J Econ Entomol* 1978; 71:397–400.
69. Lindsay IS, McAndless JM. Permethrin-treated jackets versus repellent-treated jackets and hoods for personal protection against black flies and mosquitoes. *Mosq News* 1978; 38:350–356.
70. Schreck CE, Smith N, Weidhaas D, et al. Repellents vs. toxicants as clothing treatments for protection from mosquitoes and other biting flies. *J Econ Entomol* 1978; 71:919–922.
71. Gupta RK, Rutledge LC, Reifenrath WG, et al. Effects of weathering on fabrics treated with permethrin for protection against mosquitoes. *J Am Mosq Control Assoc* 1989; 5:176–179.
72. Sholdt LL, Schreck CE, Qureshi A, et al. Field bioassays of permethrin-treated uniforms and a new extended duration repellent against mosquitoes in Pakistan. *J Am Mosq Control Assoc* 1988; 4:233–236.
73. Harback RE, Tang DB, Wirtz RA, Gingrich JB. Relative repellency of two formulations of N,N-diethyl-3-methylbenzamide (deet) and permethrin-treated clothing against *Culex sitiens* and *Aedes vigilax* in Thailand. *J Am Mosq Control Assoc* 1990; 6:641–644.
74. Eamsila C, Frances SP, Strickman D. Evaluation of permethrin-treated military uniforms for personal protection against malaria in northeastern Thailand. *J Am Mosq Control Assoc* 1994; 10:515–521.
75. Deparis X, Frere B, Lamizana M, et al. Efficacy of permethrin-treated uniforms in combination with deet topical repellent for protection of French military troops in Cote D'Ivoire. *J Med Entomol* 2004; 41:914–921.
76. Miller RJ, Wing J, Cope SE, et al. Repellency of permethrin-treated battle-dress uniforms during Operation Tandem Thrust 2001. *J Am Mosq Control Assoc* 2004; 20:462–464.
77. Khoobdel M, Shayeghi M, Vatandoost H, et al. Field evaluation of permethrin-treated uniforms against *Anopheles stephensi* and 4 species of *Culex* (Diptera: Culicidae) in Iran. *J Entomol* 2006; 3:108–118.
78. Schreck CE, Snoddy EL, Mount GA. Permethrin and repellents as clothing impregnants for protection from the Lone Star tick. *J Econ Entomol* 1980; 73:436–439.
79. Schreck CE, Mount GA, Carlson DA. Wear and wash persistence of permethrin used as clothing treatment for personal protection against the lone star tick (Acari:Ixodidae). *J Med Entomol* 1982; 19:143–146.
80. Mount GA, Snoddy EL. Pressurized sprays of permethrin and deet on clothing for personal protection against the Lone Star tick and the American dog tick (Acari:Ixodidae). *J Econ Entomol* 1983; 76:529–531.
81. Evans SR, Korch GW Jr, Lawson MA. Comparative field evaluation of permethrin and deet-treated military uniforms for personal protection against ticks (Acari). *J Med Entomol* 1990; 27:829–839.
82. Rey JL. Moyens actuels de protection contre les maladies transmises par les tiques. *Med Mal Infect* 1998; 28:393–395.
83. Ho-Pun-Cheung T, Lamarque D, Josse R, et al. Effet protecteur de vêtements imprégnés de perméthrine vis-à-vis de *D. reticulatus* et *D. marginatus* dans un biotope ouvert du centre-ouest de la France. *Bull Soc Pathol Exot* 1999; 92:337–340.
84. Faulde M, Scharninghausen J, Tisch M. Preventive effect of permethrin-impregnated clothing to *Ixodes ricinus* ticks and associated *Borrelia burgdorferi* s.l. in Germany. *Int J Med Micro* 2008; 298(Suppl 1):321–324.
85. Breeden GC, Schreck CE, Sorensen AL. Permethrin as a clothing treatment for personal protection against chigger mites (Acarina:Trombiculidae). *Am J Trop Med Hyg* 1982; 31:589–592.
86. Frances SP, Yeo AET, Burke EW, Sweeney AW. Clothing impregnations of dibutylphthalate and permethrin as protectants against a chigger mite, *Eutrombicula bisti* (Acari:Trombiculidae). *J Med Entomol* 1992; 29:907–910.
87. Mehr ZA, Rutledge LC, Inase JL. Evaluation of commercial and experimental repellents against *Xenopsylla cheopis* (Siphonaptera:Pulicidae). *J Med Entomol* 1984; 21:665–669.
88. Nassif M, Brooke JP, Hutchison DBA, et al. Studies with permethrin against body lice in Egypt. *Pestic Sci* 1980; 11:679–684.
89. Sholdt LL, Rogers EJ Jr, Gerberg EJ, Schreck CE. Effectiveness of permethrin-treated military uniforms fabric against human body lice. *Mil Med* 1989; 154:90–93.
90. Wirtz RA, Rowton ED, Hallam JA, et al. Laboratory testing of repellents against the sand fly *Phlebotomus papatasi* (Diptera:Psychodidae). *J Med Entomol* 1986; 23:64–67.
91. Asilian A, Sadeghinia A, Sharati F, et al. Efficacy of permethrin-impregnated uniforms in the prevention of cutaneous leishmaniasis in Iranian soldiers. *J Clin Pharm Ther* 2003; 28:175–178.
92. Buescher MD, Rutledge LC, Wirtz RA, Nelson JH. Laboratory repellent tests against *Rodnius prolixus* (Heteroptera:Reduviidae). *J Med Entomol* 1985; 22:49–53.
93. Kroeger A, Ordonez-Gonzalez J, Behrend M, Alvarez G. Bed net impregnation for Chagas disease control: a new perspective. *Trop Med Int Health* 1999; 4:194–198.
94. Sholdt LL, Schreck CE, Mwangiwa MI, et al. Evaluation of permethrin-impregnated clothing and three topical repellent formulations of deet against tsetse flies in Zambia. *Med Vet Entomol* 1989; 3:153–158.
95. Debboun M, Strickman DA, Klun JA. Repellents and the military: our first line of defense. *J Am Mosq Control Assoc* 2005; 21(Suppl):4.
96. Lengeler C, Snow R. From efficacy to effectiveness: insecticide treated bednets in Africa. *Bull WHO* 1996; 74:325–332.
97. Lengeler C. Insecticide-treated bed nets and curtains for preventing malaria. *Cochrane Database Syst Rev* 2004, 2. Art. No.: CD000363. DOI: 10.1002/14651858.CD000363.pub2
98. Kroeger A, Gonzalez M, Ordonez-Gonzalez J. Insecticide-treated materials for malaria control in Latin America: to use or not use? *Trans R Soc Trop Med Hyg* 1999; 3:565–570.
99. Kroeger A, Avinna A, Ordonez-Gonzalez J, Escandon C. Community cooperatives and insecticide treated materials for malaria control: a new experience in Latin America. *Malar J* 2002; 1:15.
100. Rowland M, Durrani N, Hewitt S, et al. Permethrin treated chadders and top-sheets: appropriate technology

- for protection against malaria in Afghanistan and other complex emergencies. *Trans R Soc Trop Med Hyg* 1999; 93:465–472.
101. Pennetier C, Corbel V, Hougard JM. Combination of a non-pyrethroid insecticide and a repellent: a new approach for controlling knockdown-resistant mosquitoes. *Am J Trop Med Hyg* 2005; 72:739–744.
 102. Pennetier C, Corbel V, Boko P, et al. Synergy between repellents and non-pyrethroid insecticides strongly extends the efficacy of treated nets against *Anopheles gambiae*. *Malar J* 2007; 6:38.
 103. N'Guessan R, Darriet F, Doannio JMC, et al. Olyset Net efficacy against pyrethroid-resistant *Anopheles* and *Culex* after 3 years' field use in Cote d'Ivoire. *Med Vet Entomol* 2001; 15:97–104.
 104. Kroeger A, Skovmand O, Phan QC, Boewono DT. Combined field and laboratory evaluation of a long-term impregnated bednet, PermaNet. *Trans R Soc Trop Med Hyg* 2004; 98:152–155.
 105. Lindblade KA, Dotson E, Hawley WA, et al. Evaluation of long-lasting insecticidal nets after 2 years of household use. *Trop Med Int Health* 2005; 10:1141–1150.
 106. Graham K, Kayedi MH, Maxwell C, et al. Multi-country field trials comparing wash-resistance of PermaNet and conventional insecticide-treated nets against anopheline and culicine mosquitoes. *Med Vet Entomol* 2005; 19:72–83.
 107. Kilian A, Byamukama W, Pigeon O, et al. Long-term field performance of a polyester-based long-lasting insecticidal mosquito net in rural Uganda. *Malar J* 2008; 7:49.
 108. Heal JD, Surgeoner GA, Lindsay LR. Permethrin as a tent treatment for protection against field populations of *Aedes* mosquitoes. *J Am Mosq Control Assoc* 1995; 11:99–102.
 109. Chouaibou M, Simard F, Chandre F, et al. Efficacy of bifenthrin impregnated bednets against *Anopheles funestus* and pyrethroid-resistant *Anopheles gambiae* in North Cameroon. *Malar J* 2006; 5:77.
 110. McGinn D, Frances SP, Sweeney AW, et al. Evaluation of Bistar 80SC (bifenthrin) as a tent treatment for protection against mosquitoes in Northern Territory, Australia. *J Med Entomol* 2008; 45:1087–1091.
 111. Boulware DR, Beisang AA III. Passive prophylaxis with permethrin-treated tents reduces mosquito bites among North American summer campers. *Wild Environ Med* 2005; 16:9–15.
 112. Behrens RH, Steffen R. Travel health. In: Cook GC, Zumla AI, eds. *Manson's tropical diseases*. 22nd Ed. London: Saunders Elsevier, 2009:515–525.
 113. Hewitt SE, Farhan M, Urhman N, et al. Self-protection from malaria vectors in Pakistan: an evaluation of popular existing methods and appropriate new techniques in Afghan refugee communities. *Ann Trop Med Parasitol* 1996; 90:337–344.
 114. Hewing AN, Sullivan WN, Schechter MS. Effectiveness of several pyrethroid vapors against *Aedes aegypti* (L) and *Musca domestica* L. *Mosq News* 1974; 34:54–57.
 115. Chadwick PR, Lord CJ. Tests of pyrethroid vaporising mats against *Aedes aegypti* (L) (Diptera: Culicidae). *Bull Entomol Res* 1977; 67:667–674.
 116. Warui CM. A laboratory evaluation of vaporising mats based on pyrethrins and bioallethrin for mosquito repellency. *Pyrethrum Post* 1992; 18:131–138.
 117. Amalraj DD, Kalyanasundaram M, Das PK. Evaluation of EMD vaporizers and bioallethrin vaporizing mats against mosquito vectors. *Southeast Asian J Trop Med Public Health* 1992; 23:474–478.
 118. Manga L, Robert V, Carnevale P. Efficacité des serpents et des diffuseurs en plaquettes dans la protection contre les vecteurs du paludisme au Cameroun. *Cahiers Santé* 1995; 5:85–88.
 119. Schoepke A, Steffen R, Gratz N. Effectiveness of personal protection measures against mosquito bites for malaria prophylaxis in travellers. *J Travel Med* 1998; 5:188–192.
 120. Srinivas G, Amalraj RE, Dhanraj B. The use of personal protection measures against malaria in an urban population. *Public Health* 2005; 19:415–417.
 121. Sharma VP, Nagpal BN, Srivastava A. Effectiveness of neem oil mats in repelling mosquitoes. *Trans R Soc Trop Med Hygiene* 1993; 87:626.
 122. Matsika-Claquin MD, Ménard D, Fontanet AL, et al. Efficacy of chloroquine-proguanil malaria prophylaxis in a non-immune population in Bangui, Central African Republic: a case-control study. *Trans R Soc Trop Med Hyg* 2006; 100:381–386.
 123. Liu WK, Wong MH, Miu YL. Toxic effects of mosquito coil (a mosquito repellent) smoke on rats I. Properties of the mosquito coil and its smoke. *Toxicol Lett* 1987; 39:223–230.
 124. Chen SC, Wong RH, Shiu LJ, et al. Exposure to mosquito coil smoke may be a risk factor for lung cancer in Taiwan. *J Epidemiol* 2008; 18:19–25.
 125. Liu W, Zhang J, Hashim JH, et al. Mosquito coil emissions and health implications. *Environ Health Perspect* 2003; 111:1454–1460.
 126. Heymann DL. *Control of communicable diseases manual*. 19th Ed. Washington, DC: American Public Health Association, 2008.
 127. Stürchler MP. The vector and measures against mosquito bites. In: Schlagenhauf-Lawlor P, ed. *Travelers' malaria*. 2nd Ed. Hamilton, Ontario: BC Decker, 2008:88–106.
 128. World Health Organization. *International travel and health*. Geneva: WHO, 2008.
 129. Chiodini P, Hill D, Laloo D, et al. Guidelines for malaria prevention in travellers from the United Kingdom. London: Health Protection Agency, 2007.
 130. Hill DR. Pretravel advice and immunization. In: Cohen J, Powderly WG, eds. *Infectious diseases*. 2nd Ed. London: Mosby, 2004:1429–1444.
 131. Lawrance CE, Croft AM. Do mosquito coils prevent malaria? A systematic review of trials. *J Travel Med* 2004; 11:92–96.
 132. Moore DAJ, Grant AD, Armstrong M, et al. Risk factors for malaria in UK travellers. *Trans R Soc Trop Med Hyg* 2004; 98:55–63.
 133. Hill N. Clinical evaluation of plant-based insect repellents against malaria in the Bolivian Amazon and coils in China. 13th International Congress of Entomology; 2008 July 6–12; Durban, South Africa.
 134. Lindsay LR, Surgeoner GA, Heal JD, Gallivan GJ. Evaluation of the efficacy of 3% citronella candles and 5% citronella incense for protection against field populations of *Aedes* mosquitoes. *J Am Mosq Control Assoc* 1996; 12:293–294.
 135. Müller GC, Junnila A, Kravchenko VD, et al. Indoor protection against mosquito and sand fly bites: a

- comparison between citronella, linalool and geraniol candles. *J Am Mosq Control Assoc* 2008; 24:150–153.
136. Müller GC, Junnila A, Kravchenko V, et al. Ability of essential oil candles to repel biting insects in high and low biting pressure environments. *J Am Mosq Control Assoc* 2008; 24:154–160.
 137. Coosemans M, Guillet P. La protection du voyageur contre les piqûres des moustiques. *Méd Mal Infect* 1999; 29(Suppl 3):390–396.
 138. Croft AM. Extracts from “Clinical Evidence”. *Malaria: prevention in travellers*. *BMJ* 2000; 321:154–160.
 139. Mafong EA, Kaplan LA. Insect repellents. What really works? *Postgrad Med* 1997; 102:63–74.
 140. Magnon GJ, Robert LL, Kline DL, Roberts LW. Repellency of two deet formulations and Avon Skin-so-Soft® against biting midges (Diptera: Ceratopogonidae) in Honduras. *J Am Mosq Control Assoc* 1991; 7: 80–82.
 141. Sjogren RD. An effective repellent for *Leptoconops kerteszi* Kieffer (Diptera: Ceratopogonidae). *Mosq News* 1971; 31:115–116.
 142. Asilian A, Sadeghinia A, Shariati F, et al. Efficacy of permethrin-impregnated uniforms in the prevention of cutaneous leishmaniasis in Iranian soldiers. *J Clin Pharm Ther* 2003; 28:175–178.
 143. Lindsay LR, Heal JD, Surgeoner GA. Comparative evaluation of the efficacy of Bite Blocker, OFF! Skintastic, and Avon Skin-So-Soft to protect against *Aedes* species mosquitoes in Ontario: final report. August 1996. Unpublished, 5 pp. Department of Environmental Biology, University of Guelph, Guelph, Ontario, Canada/Chemfree Environment Inc., Quebec, Canada.
 144. Proposed Regulatory Decision Document PRDD-99-02 Version, P. D. F., Statement on Personal Protective Measures to Prevent Arthropod Bites.
 145. Rajan TV, Hein M, Porte P, Wikel S. A double-blinded, placebo-controlled trial of garlic as a mosquito repellent: a preliminary study. *Med Vet Entomol* 2005; 19:84–89.
 146. Ives AR, Paskewitz SM. Testing vitamin B as a home remedy against mosquitoes. *J Am Mosq Control Assoc* 2005; 21:213–217.
 147. Hudson JE, Esozed S. The effects of smoke from mosquito coils on *Anopheles gambiae* and *Mansonia uniformis* in verandah-trap huts at Magugu, Tanzania. *Bull Entomol Res* 1971; 61:247–265.
 148. Ansari MA, Razdan RK. Relative efficacy of various oils in repelling mosquitoes. *Indian J Malariol* 1995; 32:104–111.